Inventor(s) Name: Gu, et.al.

Attorney Docket No.: 1559 WO/US

## **CLAIM AMENDMENTS**

## **IN THE CLAIMS**:

1. (Previously presented) A process for the regio- and stereoselective opening of an epoxide ring comprising:

reacting a compound having the epoxide ring with at least one amine in the presence of at least one Lewis acid.

- 2. (Currently amended.) The process according to claim 1 wherein the at least one Lewis acid is selected from the group consisting of alkyl metal halides and metal alkyl halide ethers.
- 3. (Previously presented) A process for preparing a (+)-p-mentha-2,8-diene-1-ol analog, the process comprising:

2

STLD01-1014431-1 (6/03)

Inventor(s) Name: Gu, et.al.

Attorney Docket No.: 1559 WO/US

$$H_3C$$
 $NR_1R_2$ 
 $O$ 
 $NR_1R_2$ 
 $O$ 
 $R_4$ 
 $R_4$ 

Formula (2a) Formula (4a) Formula (5a)

wherein R<sub>1</sub> and R<sub>2</sub> are H, alkyl or aryl;

wherein R<sub>4</sub> is an alkyl, alkenyl or alcohol;

wherein step (a) comprises reacting a (+)-limonene oxide analog having the Formula (1a) with at least one amine of the formula R<sub>1</sub>R<sub>2</sub>R<sub>3</sub>N in the presence of at least one Lewis acid to form amine adducts having the Formula (2a) and Formula (3a);

wherein step (b) comprises oxidizing the amine adduct of Formula (2a)to form an Noxide having Formula (4a); and wherein step (c) comprises pyrolizing the Noxide of Formula (4a) to form a (+)-p-mentha-2,8-diene-1-ol analog of Formula (5a).

- 4. (Currently amended.) The process according to claim 3 wherein the at least one Lewis acid is selected from the group consisting of alkyl metal halides and metal alkyl halide ethers.
- 5. (Previously presented) A process for preparing (+)-p-mentha-2,8-diene-1-ol, the process comprising:

Inventor(s) Name: Gu, et.al.

Attorney Docket No.: 1559 WO/US

$$\begin{array}{c} CH_3 \\ CH_3 \\ CH_3 \\ CH_3 \\ CH_3 \\ CH_3 \\ NR_1R_2 \\ OH \\ CH_3 \\ O$$

step a

Formula (1b) Formula (2b) Formula (3b)

wherein R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are H, alkyl or aryl groups;

wherein step (a) comprises reacting (+)-limonene oxide having the Formula (1b) with at least one amine of the formula  $R_1R_2R_3N$  in the presence of at least one Lewis acid to form amine adducts having the Formula (2b)and Formula (3b);

Inventor(s) Name: Gu, et.al.

Attorney Docket No.: 1559 WO/US

wherein step (b) comprises oxidizing the amine adduct of Formula (2b)to form an N-

oxide having Formula (4b); and

wherein step (c) comprises pyrolizing the N-oxide of Formula (4b) to form the (+)-p-

mentha-2,8-diene-1-ol of Formula (5b).

6. (Previously presented) The process according to claim 5 wherein the at least one amine is

selected from the group consisting of primary amines wherein R<sub>1</sub> is an alkyl or aryl group and R<sub>2</sub>

and  $R_3$  are H; secondary amines wherein  $R_1$  and  $R_2$  are alkyl or aryl groups and  $R_3$  is H; and

tertiary amines wherein R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are alkyl or aryl groups.

7. (Currently amended.) The process according to claim 5 wherein the at least one Lewis

acid is selected from the group consisting of alkyl metal halides and metal alkyl halide ethers.

8. (Cancelled.)

9. (Previously presented) The process according to claim 5 comprising oxidizing the amine

adduct of Formula (2b) by reacting the amine adduct of Formula (2b) with at least one peracid to

form the N-oxide of Formula (4b).

10. (Previously presented) The process according to claim 5 comprising oxidizing the amine

adduct of Formula (2b) by reacting the amine adduct of Formula (2b) with hydrogen peroxide in

at least one alcohol to form the N-oxide of Formula (4b).

11. (Previously presented) The process according to claim 5 comprising pyrolizing the N-

oxide of Formula (4b) in a solvent system including toluene in the presence of an at least one

particulate matter selected from the group consisting of zeolites and silica gels.

Inventor(s) Name: Gu, et.al.

Attorney Docket No.: 1559 WO/US

- 12. (Previously presented) The process according to claim 5 further including recovering the amine adduct of Formula (2b) by converting the amine adduct to an acid salt of the amine adduct of Formula (2b) by reaction with concentrated acid.
- 13. (Previously presented) A process for preparing a (+)-p-mentha-2,8-diene-1-ol analog, the process comprising:

$$CH_3$$
  $CH_3$   $CH_3$   $NR_1R_2$   $CH_3$   $NR_1R_2$   $OH$   $R_4$   $R_4$   $R_4$ 

step a

wherein R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are H, alkyl or aryl groups;

wherein R<sub>4</sub> is an alkyl, alkenyl or alcohol;

wherein R5 is an H, alkyl or aryl;

Inventor(s) Name: Gu, et.al.

Attorney Docket No.: 1559 WO/US

wherein X is a halide;

wherein step (a) comprises reacting a (+)-limonene oxide analog having the Formula (1a) with at least one amine of the formula R<sub>1</sub>R<sub>2</sub>R<sub>3</sub>N in the presence of at least one Lewis acid to form amine adducts having the Formula (2a) and Formula (3a);

wherein step (b) comprises converting the amine adduct of Formula (2a) to the acid salt of Formula (6a); and

wherein step (c) comprises base hydrolyzing Formula (6a) to form the (+)-p-mentha-2,8diene-1-ol analog of Formula (5a).

- 14. (Previously presented) The process according to claim 13 wherein the at least one amine is selected from the group consisting of primary amines wherein R<sub>1</sub> is an alkyl or aryl group and R<sub>2</sub> and R<sub>3</sub> are H; secondary amines wherein R<sub>1</sub> and R<sub>2</sub> are alkyl or aryl groups and R<sub>3</sub> is H; and tertiary amines wherein R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are alkyl or aryl groups.
- 15. (Currently amended.) The process according to claim 13 wherein the at least one Lewis acid is selected from the group consisting of alkyl metal halides and metal alkyl halide ethers.
- 16. (Previously presented) The process according to claim 13 wherein R<sub>5</sub>X is methyl iodide.
- 17. (Cancelled)
- 18. (Previously presented) A process for preparing (+)-p-mentha-2,8-diene-1-ol, the process comprising:

Inventor(s) Name: Gu, et.al.

Attorney Docket No.: 1559 WO/US

$$CH_3$$
  $CH_3$   $CH_3$ 

step a

Formula (1b)

Formula (2b)

Formula (3b)

Formula (2b)

Formula (6b)

Formula (5b)

wherein R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are H, alkyl or aryl groups;

wherein R5 is an H, alkyl or aryl;

wherein X is a halide;

Inventor(s) Name: Gu, et.al.

Attorney Docket No.: 1559 WO/US

wherein step (a) comprises reacting (+)-limonene oxide having the Formula (1b) with at least one amine of the formula R<sub>1</sub>R<sub>2</sub>R<sub>3</sub>N in the presence of at least one Lewis acid to form amine adducts having the Formula (2b) and Formula (3b);

wherein step (b) comprises converting the amine adduct of Formula (2b) to the acid salt of Formula (6b); and

wherein step (c) comprises base hydrolyzing Formula (6b) to form the (+)-p-mentha-2,8diene-1-ol of Formula (5b).

- 19. (Previously presented) The process according to claim 18 wherein the at least one amine is selected from the group consisting of primary amines wherein R<sub>1</sub> is an alkyl or aryl group and R<sub>2</sub> and R<sub>3</sub> are H; secondary amines wherein R<sub>1</sub> and R<sub>2</sub> are alkyl or aryl groups and R<sub>3</sub> is H; and tertiary amines wherein  $R_1$ ,  $R_2$  and  $R_3$  are alkyl or any groups.
- 20. (Currently amended.) The process according to claim 18 wherein the at least one Lewis acid is selected from the group consisting of alkyl metal halides and metal alkyl halide ethers.
- 21. (Previously presented) The process according to claim 18 wherein  $R_5X$  is MeI.
- 22. (Cancelled)
- 23. (Previously presented) A method for the diastereomeric separation of a mixture of (+)cis-limonene oxide and (+)-trans-limonene, the method comprising: reacting the mixture with an amine in the presence of a Lewis acid; and recovering the (+)-cis-limonene oxide that does not react with the amine.

Inventor(s) Name: Gu, et.al.

Attorney Docket No.: 1559 WO/US

24. (New) The process according to Claim 1 wherein the Lewis acid is selected from the

group consisting of lithium acetate, lithium bromide, lithium chloride, aluminum oxide and

mixtures thereof.

(New) The process according to Claim 3 wherein the Lewis acid is selected from the 25.

group consisting of lithium acetate, lithium bromide, lithium chloride, aluminum oxide and

mixtures thereof.

26. (New) The process according to Claim 5 wherein the Lewis acid is selected from the

group consisting of lithium acetate, lithium bromide, lithium chloride, aluminum oxide and

mixtures thereof.

27. (New) The process according to Claim 13 wherein the Lewis acid is selected from the

group consisting of lithium acetate, lithium bromide, lithium chloride, aluminum oxide and

mixtures thereof.

28. (New) The process according to Claim 18 wherein the Lewis acid is selected from the

group consisting of lithium acetate, lithium bromide, lithium chloride, aluminum oxide and

mixtures thereof.